



PLANTS GROWN IN UTAH THAT ARE SUSCEPTIBLE TO JAPANESE BEETLE



EXTENSIVE DAMAGE

Fruits/Vegetables

- Apple *Malus spp.*
- Asparagus *Asparagus officinalis*
- Apricot *Prunus armeniaca*
- Sweet cherry *Prunus avium*
- Sour cherry *Prunus cerasus*
- Garden plum *Prunus domestica*
- Peach *Prunus persica*
- Nectarine *Prunus persica var. nectarina*
- Japanese plum *Prunus salicina*
- Garden rhubarb *Rheum rhabonticum*
- Highbush blueberry *Vaccinium corymbosum*
- American bunch grape *Vitis labrusca*
- European grape *Vitis vinifera*

- Red raspberry *Rubus idaeus*
- Corn *Zea mays*
- Soybean *Glycine max*
- Basil *Ocimum basilicum*

Ornamentals

- Japanese maple *Acer palmatum*
- Norway maple *Acer platanoides*
- Horse chestnut *Aesculus hippocastanum*
- Hollyhock *Althaea rosea*
- Rose of Sharon *Hibiscus syriacus*
- Black walnut *Juglans nigra*
- Japanese kerria *Kerria japonica*
- Crabapple *Malus baccata*
- Japanese flowering crabapple *Malus floribunda*
- Evening primrose *Oenothera biennis*

- Virginia-creeper *Parthenocissus quinquefolia*
- London planetree *Platanus x acerifolia*
- Lombardy poplar *Populus nigra Italica*
- Rose *Rosa spp.*
- Purple leaf plum *Prunus cerasifera*
- Silver lace *Polygonum (Fallopia) aubertii*
- Whirling Butterflies *Gaura spp.*
- Peking cotoneaster *Cotoneaster acutifolius*
- Cinquefoil *Potentilla spp.*
- Pussy willow *Salix discolor*
- American mountain-ash *Sorbus americana*
- American linden *Tilia americana*
- American elm *Ulmus americana*
- Turf (various genera)
- Industrial/Medicinal**
- Hemp/marijuana *Cannabis sativa*

MODERATE DAMAGE

Fruits/Vegetables

- Cauliflower *Brassica oleracea var. botrytis*
- Garden bean *Phaseolus vulgaris*
- Black raspberry *Rubus occidentalis*
- Garden red currant *Ribes sativum*

Ornamentals

- European black alder *Alnus glutinosa*
- European white birch *Betula pendula*
- Southern catalpa *Catalpa bignonioides*
- Garden canna *Canna generalis*
- Dahlia *Dahlia variabilis*
- Marigold *Tagetes spp.*
- Common zinnia *Zinnia elegans*

- Pin oak *Quercus palustris*
- Chinese wisteria *Wisteria sinensis*
- Butterflybush *Buddleja davidii*
- Common four-o'clock *Mirabilis jalapa*
- European larch *Larix decidua*
- Baldcypress *Taxodium distichum*
- American planetree *Platanus occidentalis*
- Common quince *Cydonia oblonga*
- Oriental cherry *Prunus serrulata*
- Babylon weeping willow *Salix babylonica*
- European linden *Tilia cordata*
- Cattail *Typha latifolia*
- Livestock Feed**
- Alfalfa *Medicago sativa*
- Alsike clover *Trifolium hybridum*
- Red clover *Trifolium pratense*



References:
 Cranshaw, W. 2018. Japanese Beetle. Colorado State University Extension Fact Sheet 5.601.
 Fleming, W.E. 1972. Biology of the Japanese beetle. Technical Bulletin 1449 of the United States Department of Agriculture. Washington, DC. 129 pp.
 University of Minnesota Extension. <https://extension.umn.edu/yard-and-garden-insects/japanese-beetles> [visited on web 10 Feb. 2020]
 Williamson, C.R. 2013. Japanese Beetle. University of Wisconsin-Madison Fact Sheet XHT 1062.

Photo credits:
 Top (left to right): 1) Whitney Cranshaw, Colorado State University 2) M.J. Klein, USDA ARS 3) Bruce Watt, University of Maine.
 Bottom (left to right): 1) Sarah Schulthies, Utah Department of Agriculture and Food 2) Daren Mueller, Iowa State University 3) Steven Katovich, Bugwood.org 4) White Cranshaw, Colorado State University.

Thanks to Salt Lake County USU Extension for reviewing this document.
 "Plants grown in Utah that are susceptible to Japanese beetle" poster V1.0
 Utah Department of Agriculture and Food—Plant Industry 2020