

Badge X2

Fungicide/Bactericide

FOR ORGANIC PRODUCTION



EPA Reg. No. 80289-12-10163

For the control of Cryptogams on ornamentals in greenhouses, shadehouses, outdoor nurseries and outdoor landscape planting in the State of Alabama, Arizona, Arkansas, California, Colorado, Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maine, Maryland, Massachusetts, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, Nevada, New Hampshire, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, South Dakota, Tennessee, Texas, Utah, Vermont, Virginia, Washington, West Virginia, Wisconsin, Wyoming.

FIFRA §2(ee) RECOMMENDATION: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the United States Environmental Protection Agency (EPA).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product bulletin must be in the possession of the user at the time of pesticide application of Badge® X2 Fungicide/Bactericide. This labeling contains supplemental instructions that may not appear on the product label. All applicable directions, restrictions, and precautions on the product label are to be followed.

CROP	DISEASE	RATE (LBS PER ACRE)	MAX. APP. RATE	USE INSTRUCTIONS
Ornamentals (specified as listed on the product label)	Cryptogams, such as algae, nostoc, lichens, and moss	1.5-5	Unless otherwise noted, Single: 2 lbs Cu/A Annual: 20 lbs Cu/A	For ornamental crops in dormancy, apply as a thorough cover spray at rates ranging from 1.5 to 5 lbs/A of BADGE X2. When new growth is present, apply as a thorough cover spray at rates ranging from 1.5 to 2 lbs/A of BADGE X2. One and a half (1.5) level tablespoons of BADGE X2 per 1000 square feet is equivalent to 2.4 lbs/A. Begin application at first sign of disease and repeat at 7 to 14 day intervals; use the higher rates and shorter spray intervals during periods of frequent rains or when severe disease conditions persist. The minimum retreatment interval is 7 days.
<ul style="list-style-type: none"> Plant sensitivities to BADGE SC have been found to be acceptable for the specific genera and species listed on the product label under the conditions tested. However, phytotoxicity may occur. Due to the large number of species and varieties of ornamental and nursery plants and the wide range of growing conditions, it is impossible to test every one for sensitivity to BADGE SC. Neither the manufacturer nor the seller has determined whether or not BADGE SC can be safely used on ornamental or nursery plants not listed on the product label. The user should determine if BADGE SC can be used safely prior to commercial use. In a small area, apply the labeled rates to the plants in question, (bedding plants, foliage, etc.), and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use. This product may be reactive on masonry and metal surfaces such as galvanized roofing. Avoid contact with metal surfaces. Do not spray on cars, houses, lawn furniture, or other metallic surfaces. 				

Badge is a registered trademark of Isagro
ID# 19-10-04



Manufactured by Isagro SpA for:
Isagro USA, Inc.,
430 Davis Drive, Suite 240,
Morrisville, NC 27560



Distributed by :
Gowan Company LLC
P.O. Box 5569
Yuma, AZ 85366-5569