



## FIFRA 2(ee) Recommendation

EPA Reg. No.: 279-9658

**FOR DISTRIBUTION AND USE ONLY  
IN REGISTERED STATES  
(EXCEPT TEXAS)**

**THIS RECOMMENDATION, WHICH CONTAINS  
ADDITIONAL DIRECTIONS FOR USE, IS MADE AS  
PERMITTED UNDER FIFRA SECTION 2(ee) AND  
HAS NOT BEEN SUBMITTED TO OR APPROVED  
BY THE US EPA.**

This recommendation is valid until December 31, 2025 or until withdrawn, canceled or suspended.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.**

**THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

**For “2x2” Placement of Xyway™ LFR at-plant in Corn (Field, Corn Grown for Seed, Popcorn, Sweet Corn)**

### Directions for Use

- Xyway LFR fungicide may be applied at planting time to the soil in-furrow with the seed or placed sub-surface off the row. (i.e. 2x2 inch, 2x2x2 dual-placement, or similar placements) within 3-inches of the seed.
- For control of late-season infestations, heavy disease pressure situations or foliar diseases not listed, a supplemental foliar application may be needed.
- When Xyway LFR placement is away from the seed row in a 2x2, a 2x2x2 or similar dual placement, apply full use rate (15.2 fl. oz of product/A) for best results.
- Refer to row space conversion chart on the EPA registered Xyway LFR fungicide label for application rates on row spacings other than 30”.

### Restrictions:

- **Do not apply more than 0.228 lb. active ingredient per acre (15.2 fl oz of product/A) per year including at-plant plus foliar applications of other flutriafol-containing products.**
- **Do not apply this product as a foliar application.**

R-4520 012621 11-04-20



FMC Corporation  
2929 Walnut St.  
Philadelphia, PA 19104  
215-299-6000