



FIFRA 2(ee) Recommendation
EPA Reg. No.: 279-9658
FOR DISTRIBUTION AND USE ONLY IN REGISTERED STATES (EXCEPT TEXAS)
THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE US EPA.

This recommendation is valid until December 31, 2025 or until withdrawn, canceled or suspended. Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.
THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

For Suppression of Anthracnose Stalk Rot, Fusarium Stalk and Crown Rot and Physoderma Brown Spot in Corn (Field, Corn Grown for Seed, Popcorn)

Directions for Use
Apply 0.87 fl. oz./1000 row-ft. Xyway™ LFR fungicide (15.2 fl. oz./A on 30" rows) at-planting for suppression of anthracnose stalk rot, stalk and crown rot diseases caused by susceptible <i>Fusarium</i> spp., and <i>Physoderma</i> brown spot.

CROP	PEST	RATE OF APPLICATION		
		Fluid oz/A (30" row spacing)	Fluid oz/1000 row-feet	Lbs. A.I./acre
CORN (Field, Corn Grown for Seed, Popcorn)	Anthracnose Stalk Rot (<i>Colletotrichum graminicola</i>) Fusarium Stalk and Crown Rot (<i>Fusarium graminearum</i> , <i>Fusarium</i> spp.) Physoderma Brown Spot (<i>Physoderma maydis</i>)	15.2	0.87	0.228

Restrictions:

- **Do not apply more than 0.228 lb. active ingredient per acre (15.2 fl oz of product/A) per year including at-plant plus foliar applications of other flutriafol-containing products.**
- **Do not apply this product as a foliar application.**

R-4521 012621 11-04-20



FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000