


# AMERICAN FOULBROOD

PREVENTION  
DIAGNOSIS  
CONTROL



American foulbrood (AFB) is the most devastating of the honey bee brood diseases. It is easily transmittable and highly deadly. The condition is caused by the spore-forming bacteria *Paenibacillus larvae*<sup>1,2</sup>. Just a small number of spores can infect healthy three-day old larvae<sup>3</sup>. While the bacterium can be killed by antimicrobials or environmental extremes<sup>4</sup>, the millions of spores it produces are extremely resilient and can remain viable in honey and beekeeping equipment for over 40 years<sup>5,6</sup>.

## Spread of the Disease

The disease is transmitted in various ways:

- In infected colonies, young worker honey bees inadvertently spread the spores throughout the hive when they attempt to remove spore-laden dead larvae<sup>7</sup>.
- Robber bees from uninfected hives may take the pathogen back to their hives when they forage honey from a colony that is infected with the disease<sup>8</sup>.
- Humans can transmit the disease by exposing a healthy colony to contaminated bees or equipment such as frames, boxes, or tools<sup>15</sup>.

## Prevention

Preventing a hive from becoming infected is the easiest and best method for managing the disease:

- When purchasing a colony, find out if it was treated with antibiotics. Medicines may be masking symptoms of an already infected colony; the hive will likely become symptomatic if treatment ends<sup>11,16</sup>.
- Before purchasing or accepting used beekeeping equipment contact a bee inspector to have it examined for the disease.
- Sterilize tools and personal protective equipment after working in a hive you suspect is infected.
- Never switch frames, boxes, or other equipment from an infected hive to a hive that is free from disease.
- Don't share unsterilized tools or equipment with other beekeepers.
- Put unused beekeeping equipment in a bee-proof location.
- Antibiotics can be used preventatively in specific high-risk circumstances, but only under the direction of a licensed veterinarian.

## KNOW THE SYMPTOMS




Figure 1




Figure 2




Figure 3

In order to make a proper diagnosis, be familiar with signs of the disease:  
*Fig. 1:* Prepupae form dark "scales" in their cells, which the bees have difficulty removing. *Fig. 2:* Cell caps may appear sunken and dark in color and "ropy" when penetrated. *Fig. 3:* Spotty brood pattern with small perforations appear in the capping. Other symptoms include dead brood that are tan to dark brown, pupae found with extruded tongues, and a distinct, foul odor. These visual symptoms are helpful in diagnosis, however other brood maladies have similar symptoms. Testing dead brood is the best way to know if a colony is infected. Testing is available through your bee inspector.

## Control

Destruction of beekeeping equipment by burning or deep burial is the safest and most effective way to control the disease. However if the disease is detected early enough, a beekeeper may want to pursue treatment options.

- Oxytetracycline, Tylosin Tartrate, and Lincomycin are the only antibiotics that are approved for treatment of AFB<sup>17</sup>.
- To obtain antibiotics, a veterinarian must either write a Veterinary Feed Directive (VFD) or prescription. A list of veterinarians that will work with beekeepers can be found at the UDAF web address below.
- Some strains of AFB have become resistant to antibiotics<sup>12,13,14</sup>.
- If treatment is unsuccessful, burning or deep burial of the equipment is the most prudent course of action. This is necessary due to the long-lived infectious spores that are left behind on exposed equipment.



## Contact Information and Resources

This informational factsheet is brought to you by the Utah Department of Agriculture and Food.

To request a free hive health inspection call 801-982-2313 or email [udaf-insects@utah.gov](mailto:udaf-insects@utah.gov)

Find veterinarians that can prescribe antibiotics at: <https://ag.utah.gov/farmers/plants-industry/apiary-inspection-and-beekeeping/find-an-apiary-veterinarian/>

## Sources

1. White, G. F. 1920. American foulbrood. U.S. Dept of Agriculture, Washington, DC.
2. Genersch, E. 2010. American Foulbrood in honeybees and its causative agent, *Paenibacillus larvae*. *J. Invertebr. Pathol.* 103: S10–S19.
3. Brødsgaard, C. J., Ritter, W., & Hansen, H. (1998). Response of in vitro reared honey bee larvae to various doses of *Paenibacillus larvae larvae* spores. *Apidologie*, 29(6), 569-578. <https://doi.org/10.1051/apido:19980609>
4. Dobbelaere, W., De Graaf, D. C., Reybroeck, W., Desmedt, E., Peeters, J. E., & Jacobs, F. J. (2001, March 2). Disinfection of wooden structures contaminated with *Paenibacillus larvae* subsp. *larvae* spores. *Journal of Applied Microbiology*, 91, 212-216. <https://doi.org/10.1046/j.1365-2672.2001.01376.x>
5. Haseman L. (1961) How long can spores of American foulbrood live, *Am. Bee J.* 101, 298-299.
6. Forsgren, E., Stevanovic J., and Fries I. 2008. Variability in germination and in temperature and storage resistance among *Paenibacillus larvae* genotypes. *Vet. Microbiol.* 129: 342–349.
7. Lindström, A., Korpela, S., & Fries, I. (2008). The distribution of *Paenibacillus larvae* spores in adult bees and honey and larval mortality, following the addition of American foulbrood diseased brood or spore-contaminated honey in honey bee (*Apis mellifera*) colonies. *Journal of Invertebrate Pathology*, 99(1), 82-86. <https://doi.org/10.1016/j.jip.2008.06.010>
8. Lindström, A., Korpela, S., & Fries, I. (2008, October 28). Horizontal transmission of *Paenibacillus larvae* spores between honey bee (*Apis mellifera*) colonies through robbing. *Apidologie*, 39(5), 515-522. <https://doi.org/10.1051/apido:2008032>
9. Brady, S. T., Roll, C. R., Walker, J. K., Fajardo, C. P., Breakwell, D. P., Eggett, D. L., & Hope, S. (2021). Phages Bind to Vegetative and Spore Forms of *Paenibacillus larvae* and to Vegetative *Brevibacillus laterosporus*. *Frontiers in Microbiology*, 12, 21. <https://doi.org/10.3389/fmicb.2021.588035>
10. Brady, S. T., Merrill, B. D., Hilton, J. A., Payne, A. M., Stephenson, M. B., & Hope, S. (2017). Bacteriophages as an alternative to conventional antibiotic use for the prevention or treatment of *Paenibacillus larvae* in honeybee hives. *Journal of Invertebrate Pathology*, 150, 94-100. <https://doi.org/10.1016/j.jip.2017.09.010>
11. Adriana M Alippi, Graciela N Albo, Daniel Leniz, Inés Rivera, Marta L Zanelli & Amelia E Roca (1999) Comparative study of tylosin, erythromycin and oxytetracycline to control American foulbrood of honey bees, *Journal of Apicultural Research*, 38:3-4, 149-158, DOI: 10.1080/00218839.1999.11101005
12. Alippi, A. M., López, A. C., Reynaldi, F. J., Grasso, D. H., & Aguilar, M. O. (2007, December 15). Evidence for plasmid-mediated tetracycline resistance in *Paenibacillus larvae*, the causal agent of American Foulbrood (AFB) disease in honeybees. *Veterinary Microbiology*, 125(3-4), 290-303. <https://doi.org/10.1016/j.vetmic.2007.05.018>
13. Miyagi, T., Peng, C. Y.S., Chuang, R. Y., Mussen, E. C., Spivak, M. S., & Doi, R. H. (2000). Verification of oxytetracycline-resistant American foulbrood pathogen *Paenibacillus larvae* in the United States. *Journal of Invertebrate Pathology*, 75(1), 99-96. <https://doi.org/10.1006/jipa.1999.4888>
14. Evans, J. D. (2003, May). Diverse origins of tetracycline resistance in the honey bee bacterial pathogen *Paenibacillus larvae*. *Journal of Invertebrate Pathology*, 83(1), 46-50. [https://doi.org/10.1016/S0022-2011\(03\)00039-9](https://doi.org/10.1016/S0022-2011(03)00039-9)
15. Sturtevant, A. P. 1932. Relation of commercial honey to the spread of American Foulbrood. *J. Agric. Res.* 45: 257.
16. Marco Lodesani & Cecilia Costa (2005) Limits of chemotherapy in beekeeping: development of resistance and the problem of residues, *Bee World*, 86:4, 102-109, DOI: 10.1080/0005772X.2005.11417324
17. US Food and Drug Administration. (2021, August 24). FDA Approved Animal Drug Products SEARCH:BEES. Animal Drugs @ FDA. Retrieved August 24, 2021, from <https://animaldrugatfda.fda.gov/adafda/views/#/search>